



evercare

HOSPITAL DHAKA
Transforming Healthcare



Organization Accredited
by Joint Commission International

DISCHARGE SUMMARY

INJ MOXIFLOXACIN

INJ MEROPENEM

INJ LINEZOLIDE

INJ COTSON

MEDICATION ADVICE ON DISCHARGE:

TAB MOXIFLOXACIN - 400 MG- 0+0+1- AFTER MEAL- FOR 5 DAYS

TAB DOCOPA - 200 MG- 1+0+1- 2 WEEKS.

TAB PREDIXA - 16 MG- 1+1+0 - 5 DAYS AND THEN- 1+0+0 FOR 5 DAYS

TAB DICALTRDL - 0.25 MG- 0+0+1- CONTINUE ✓

TAB ECOSPRIN- 75 MG- 0+1+0- AFTER MEAL- CONTINUE ✓

CAP SECLO - 20 MG- 1+0+1- BEFORE MEAL- 2 WEEKS

TAB THYROX- 50 MCG- 3+0+0- BEFORE BREAKFAST- CONTINUE ✓

TAB CALCIUM+ VIT. D3- 500 MG/200 IU- 2+0+0- CONTINUE

TAB BETALOC- 50 MG- 1+0+0- CONTINUE ✓

TAB PRAZOPRESS- 5 MG- 1+0+1- CONTINUE ✓

TAB TELMISARTAN+HYDROCHLORTHIAZIDE- 80/12.5 MG- 1+0+0- CONTINUE SYP MUCOLYT- 2 TSF- 8 HOURLY- 1 FILE

DIET ADVICE:

LOW SALT LOW FAT RENAL DIET

OTHER ADVICE:

1. TAKE YOUR MEDICATIONS REGULARLY.