

## **DEPARTMENT OF PATHOLOGY**

## ASH-QRF-HP-07

Patient Name	Mr. MOHAMMED DALIUR RAHMAN		A	<b>\ge</b> 44Yr	Gender	Male
UHID	AC01.0004039730	Lab No ASH0	1.H19121	71 <b>LRN</b>	7818887	
W/BNo/RefNo	5th A WARD/1507A	Received on	22-OCT-2019 10:14:46 AM			
Collected on	22-OCT-2019 10:14:46 AM	PatSerNo	erNo CSNIP87798			
Reported on	26-OCT-2019 02:30:14 PM	UHID				
Ref Doctor	Dr.RECNAC - SRIPRAKASH DURAISAMY					

lymphoplasmacytic infiltrate. The tumor infiltrates buccinator muscle fibres, subcutis and papillary dermis to a depth of 2.1 cm. There is no evidence of ulceration of epidermis. Medially, the tumor does not involve the mandibular and maxillary periosteum.

H1 to H9: Sections show mucosal margins including the upper amnd lower mucocutaneous anterior margin lined by squamous epithelium with focal ulceration, free of tumor.

J1 to J3: Sections show skin lined by squamous epithelium, free of tumor.

K1 to K6: Sections show fibroadipose tissue and skeletal muscle bundles, free of tumor.

M: Sections show 2 lymph nodes with reactive hyperplasia, free of tumor.

N: Sections show 4 lymph nodes with reactive hyperplasia, free of tumor.

P: Sections show salivary gland tissue with no specific pathology.

Q1, Q2: Sections show 5 lymph nodes with reactive hyperplasia, free of tumor.

R1, R2: Sections show 1/10 lymph nodes with metastatic carcinoma with a maximum tumor deposit size of 0.3 cm and no extranodal extension.

## IMPRESSION:

PROCEDURE: WIDE LOCAL EXCISION OF ULCER RIGHT BUCCAL MUCOSA WITH CHEEK, MARGINAL MANDIBULECTOMY AND SUPERIOR ALVEOLECTOMY

HISTOLOGIC TYPE: SQUAMOUS CELL CARCINOMA, GRADE 2

TUMOR SITE: RIGHT BUCCAL MUCOSA



Page 6 of 8

Kindly bring your records during your next visit to our hospital