



DISCHARGE SUMMARY

Name of Patient	Mr. KISHORE SHARMA	Date of admission	16/04/2023
Age	60 Years	Date of discharge	27/04/2023
UHID No.	326534	Consultant: DR. MOHAMMAD HAROON-OR-RASHID MBBS, FCPS, MRCS, MS	
IP No.	23/4928		
Bed No.	605-A		
DIAGNOSIS	Carcinoma splenic flexure of colon with intestinal obstruction		

CASE SUMMARY

Mr. KISHORE SHARMA, 60 years old male, was admitted in this hospital on 16.04.2023 with complaints of abdominal pain for 2 months, H/O weight loss for 1.5 months, vomiting for 2 weeks with associated nausea. After proper evaluation and colonoscopy procedure, malignant colonic growth with luminal obstruction is found. After pre-operative preparation, Left Hemicolectomy with double barrel colostomy was done on 19.04.2023 under G/A. His Post-operative period was uneventful. Now his condition is improved and is being discharged with advice.

All Documents: Supplied

OT Note:

Date : 19.04.2023
Operation : Left hemicolectomy with double barrel colostomy
Indication : Carcinoma splenic flexure of colon with intestinal obstruction
Anesthesia : G/A
Surgeon : Dr. Md. Haroon-Or-Rashid
Anesthetist : Dr. Sanaul
Procedure : With all aseptic precautions, after proper painting & draping, abdomen opened by midline incision. Medial to lateral dissection was done. Sigmoid & descending colon mobilized from lateral wall. Left colic artery(LCA) & Inferior mesenteric vein(IMV) ligated. Medial dissection done from retro peritoneum. (Left) Ureter & gonadal vessel identified. Splenic flexure was mobilized. (Left) Branch of middle colic artery ligated. Left Hemicolectomy with double barrel colostomy done. 500ml of ascitic fluid aspirated. No peritoneal or liver metastasis. No pelvic deposit. Growth of splenic flexure of colon. Transverse colon & right colon hugely dilated. After hemostasis abdomen is closed in layers. Specimen were sent for histopathology.

Medication:

SL.	Medicine	Dose	Frequency	Duration
✓1.	Cap. Cef-3	400mg	1+0+1(After Meal)	7 days ✓
✓2.	Cap. Sergel	20mg	1+0+1 (Before Meal)	14 days ✓
✓3.	Tab. Napā	500mg	1+1+1 (After Meal)	5 days ✓
✓4.	Tab. monas	10 mg	0+0+1	15 days ✓
✓5.	Tab Ceevit	250mg	1+0+1	15 days ✓
✓6.	Tab. Omidon	10 mg	1+1+1	15 days ✓