



CENTRAL HOSPITAL LIMITED

House # 2, Road # 5, Green Road, Dhanmondi, Dhaka-1205, Bangladesh
Tel : +88-02-41060800-19, E-mail : chl@bol-online.com
info@centralhospitaltdbd.com, Web : www.centralhospitaltdbd.com



Crossmatching & Screening Report



Invoice No : I2302030450 Invoice Date : 12/02/23 Delivery Date : 12/02/23 Report No : 12302036531
Patient Name : SHAHANAZ AKTER (LOVELY) Age : 65Y Gender : F
Donor Info. : MD. RASHEDUL Age : 33Y Gender : MALE
Referred By : Dr. Indrajit Kumar Kundu
Admission No : A2301003267 Ward : CABIN - 908 (AC) Bed : CABIN - 908 (AC)

Sample : BLOOD LAB. No : 12302041769

Tests : Blood Crossmatching and Screening

Patient's Blood Group (ABO) : "B" Positive
Donor Blood Group (ABO) : "B" Positive
Type of Donation : Directed
Type of Blood Component : Whole Blood
Cross Match Status at RT & 37°C ICT method : COMPATIBLE
Bag No : 818

S.T. Given 12.02.23

Donor Screening Tests:

1. HBs Ag : Negative
2. HCV : Negative
3. HIV (1+2) : Negative
4. MP : Negative
5. Syphilis : Negative

Note : HBsAg, HCV, Syphilis, HIV (1+2) and Malarial Parasite Screening Carried out by Immunochromatographic assay method

The blood/component is ready for supply/supplied for transfusion

Note :

Instructions for use Blood Component / Product :

1. Please check the label on the blood bag, requisition form and money receipt properly before starting transfusion.
2. Use only Standard blood transfusion set, Filter for transfusion of blood and blood component.
3. Do not add any medication to the unit of blood / blood components or products.
4. Check the unit of Blood /Blood components or products for color, clot formation, leakage. If anything abnormal found please contact to blood bank.
5. Transfusion should be initiated within 30 mins of taking the blood out of the refrigerator. Once started, the transfusion should be completed within 2-3 hours. If not required please return the blood bag within 30 mins to blood bank with reason and name of the doctor. Platelet and plasma products (thawed) should be transfused as soon as possible to get optimum benefit. Thawed plasma would never be refrozen.
6. Do not warm blood bag. Warming of blood is required only in case of special circumstances. Blood bag with inserted transfusion set once used for transfusion would not be stored in blood bank for further transfusion
7. If any adverse reaction occurs stop the transfusion immediately and send the blood bag, BT set, post transfusion blood sample (10ml) from patient in EDTA and plain tubes with documentation of symptoms in a form filled by doctor.

Remarks :